SUICIDE RISK SCREENING PATHWAY **OUTPATIENT PRIMARY CARE** & SPECIALITY CLINICS **Presentation to Outpatient** Primary Care & Speciality Clinics Medically able to answer questions? NO Screen all patients ages 10 above who meet any of the screening criteria.* Screen at YES next visit *SCREENING CRITERIA Administer ASQ (ideally separate from parents) 1. New patient 2. Existing patient who has not been screened within the past 30 days 3. Patient had a positive suicide risk screen the last time they were screened YES on any question 1-4 or refuses to answer? **NEGATIVE SCREEN** 4. Clinical judgement dictates screening Exit Pathway YES If patient answered "yes" to Q4, and the patient has been screened before, ask: "Since last visit, have you tried to kill yourself?" If they answer "no" and they also answered "no' to Q1-3, no further action needed. YES YES to Q5? If the only "yes" answer is to Q4 (past suicidal behavior), factors to consider: ☐ Was the attempt more than a year ago? NO Has the patient received or is currently in mental health care? Is parent aware of past suicidal behavior? Is the suicidal behavior not a current, active concern? If yes to all these, then consider "Low Risk" choice for action Non-acute Positive Screen; Conduct Brief Suicide Safety Assessment (BSSA) Detailed instructions about the BSSA can be found at www.nimh.nih.gov/ASQ BSSA outcome(three possibilities) **IMMINENT RISK LOW RISK FURTHER EVALUATION NEEDED** Patient has acute suicidal thoughts and needs an urgent full mental health evaluation No further evaluation Mental health referral needed as soon as possible needed at this time **INITIATE SAFETY PRECAUTIONS¹** Make a safety plan with the patient and parent/guardian to activate as needed. Until able to obtain full Would benefit from mental health evaluation a non-urgent mental health If mental health evaluation follow-up? is not available within practice, refer to outpatient mental health clinician. SAFETY PRECAUTIONS Per institution protocol; keep patient under direct YES observation, remove NO dangerous items, provide safety education, etc. **REFERRAL** Schedule a follow up with patient within 72 hours for safety check and to determine whether or not they were able to obtain a mental health appointment to further mental health No referral care as appropriate; Continue needed medical care; Initiate safety at this time plan for potential future suicidal thoughts Send to emergency department for full mental health/safety evaluation SAFETY PLANNING · Create safety plan for potential future suicidal thoughts, including identifying personal warning signs, coping strategies, social contacts for support, and emergency contacts. Detailed instructions about safety planning can be found at https://www.sprc.org/resources-programs/patient-safety-plan-template • Discuss lethal means safe storage and/or removal with both parent/guardian and child (e.g. ropes, pills, firearms, belts, knives)

- Provide Resources: 24/7 National Suicide Prevention Lifeline
- 1-800-273-TALK (8255), En Español:1-888-628-9454, 24/7 Crisis Text Line: Text "START" to 741-741

If suicide risk becomes more acute, instruct patient/parent/guardian to contact outpatient healthcare provider to evaluate need for ED visit.

Schedule all patients who screen positive for a follow-up visit in 3 days to confirm safety and determine if a mental health care connection has been made.

Future follow-up primary care appointments should include re-screening patient, reviewing use of safety plan, and assuring connection with mental health clinician.